

Prescription Treatment® brand

2% Propoxur Bait

For effective control of cockroaches, crickets and ants

KILLS:

Cockroaches, Crickets and Ants

FOR USE IN AND AROUND:

Apartment Buildings, Government Offices, Homes, Institutions, Poultry Packing†, Public Buildings, Red Meat Packing and Food Processing Plants† (including plants that are federally inspected), Restaurants† (food and non-food areas), Stores† (including food processing areas), and Warehouses

† See special instructions for these sites under Directions for Use.

ACTIVE INGREDIENT:

Propoxur 2.0%

INERT INGREDIENTS: 98.0%

TOTAL: 100.0%

EPA Reg. No. 499-518

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 min. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15 - 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: Product contains a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate in large therapeutic doses is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. **SYMPTOMS OF POISONING:** A sense of "lightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT -

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS - Users

should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and aquatic invertebrates. Birds and other small mammals feeding on treated bait may be killed. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Prescription Treatment® brand 2% Propoxur Bait, a ready-to-use product, is very effective for the control of ants (except California), crickets and the cockroaches listed below. It gives rapid kill of these insects including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. Crickets and roaches die quickly after feeding on the bait. Apply as needed to control the target pests.

- Cockroaches: American, Brown, Brown banded, German, and Oriental roaches.
- Crickets
- Ants (including carpenter ants) (except in California)

USE RESTRICTIONS

- Do not formulate this product into other end use products.
- Do not apply to floors or other surfaces likely to be contacted by children or pets.
- Do not apply this product in a way that will contact any person or pet, either directly or indirectly. Keep people and pets out of the application area during application.
- Do not apply as a landscape treatment (to lawns, shrubs or trees, garden plants).

INDOOR USE

For use in Apartment Buildings, Government Offices, Homes, Institutions, Public Buildings, Restaurants (non-food), Stores and Warehouses.

Apply bait lightly to floors near baseboards, in closets, under sinks and refrigerators, around and inside



**WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.**

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information. For automatic specimen label updates, register at www.wmmg.com.



2% Propoxur Bait

garbage cans, and in other areas where insects are found but which are inaccessible to children such as cracks and crevices. (Do not make Crack & Crevice® applications in California.) Do not apply to animals. Do not treat portions of buildings where animals can feed on bait. Use approximately 4 oz of bait/1,000 ft² of space and scatter lightly so that bait is barely visible. Repeat as necessary. Keep out of reach of children.

FOOD HANDLING ESTABLISHMENTS*

For use by pest management professionals (PMPs) in Restaurants (food areas), Poultry Packing, Red Meat Packing and Food Processing Plants (including plants that are Federally inspected) Food Processing Areas of Stores, Warehouses, Government Offices, Public Buildings and Institutions.

Apply bait on paper, pasteboard, or other material that will permit removal. Locate these makeshift bait stations on window sills, on floor near walls, in storage areas and other areas where insects have been observed. Maintain strict account of location and number of bait stations. Do not apply bait in edible product areas until operations have ceased for the day. Remove all bait stations from the edible products areas before resuming operations.

Before applications are to be made in food processing plants that are inspected under the Meat and Poultry Inspection Program of USDA, a blue dye must be mixed as follows with the bait before entering the plant and setting out the bait stations:

(1) To each 4 lbs container of bait add 6 tsp of a blue food color dye (FD&C Blue No. 1).

(2) Thoroughly mix by shaking or mechanical means until the dye is uniformly distributed throughout the bait.

Do not store product in or around such processing plants.

RESEARCH FACILITIES

The use of this product in research animal quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals. Bait should not be applied directly on animals nor applied where animals can feed on the bait.

OUTDOOR USE

Use only around patios, driveways, sidewalks and foundations. Apply bait evenly in a band 2 - 3 ft wide at the rate of 4 oz/1,000 ft² around foundations, patios, driveways, or sidewalks, taking care to treat any cracks or crevices. For maximum effectiveness, treat beneath concrete drain splash pans.

STORAGE AND DISPOSAL

Do not puncture or incinerate! Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If container is partly filled, contact your local solid waste agency for disposal instructions.

A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.

3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

www.wmmg.com

© 2006 Whitmire Micro-Gen Research Laboratories, Inc.