

PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals

CAUTION: Keep away from children, domestic animals and pets. If swallowed, this product reduces the clotting ability of blood and causes bleeding. Avoid contact with eyes, skin or clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking.

FIRSTAID

Have this label with you when obtaining treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Note to Physician or Veterinarian: Contains difethialone, an anticoagulant. For humans or dogs that have ingested this product, or have obvious poisoning symptoms (bleeding) or lowered prothrombin times, give Vitamin K₁ by intramuscular or oral administration. Check prothrombin time every 3 days until values return to normal.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of empty container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg.No. 7173-236
EPA Est.No. 7173-WI-1

Product No.
Label No.



MINI BLOCKS

**KILLS NORWAY RATS,
ROOF RATS, HOUSE MICE AND
WARFARIN-RESISTANT HOUSE MICE**

**NORWAY RATS AND HOUSE MICE CAN CONSUME
A LETHAL DOSE IN ONE NIGHT'S FEEDING
WITH FIRST DEAD RODENTS APPEARING
FOUR OR FIVE DAYS AFTER FEEDING BEGINS**

Active Ingredient:difethialone* 0.0025%
Inert Ingredients 99.9975%
Total 100.0000%

*Patented in the U.S.A.

KEEP OUT OF REACH OF CHILDREN
CAUTION: See side panel for additional precautionary statements.



Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(414) 351-1476

NET WEIGHT: 15 lbs (6.8 kg)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control Norway rats, roof rats and house mice. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Urban Areas: May be used indoors and in transport vehicles (ships, trains, aircraft). May be used outdoors within 15 feet of homes, food processing facilities, industrial, commercial, agricultural and public buildings, and similar manmade structures.

Non-urban Areas: This product may only be used inside of homes and agricultural buildings.

Selection of Treatment Areas: Determine areas where rats or mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls, or in locations where rodents or their signs have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Mice: Apply one block per placement. Two blocks may be needed at points of very high mouse activity. Space placements at 8 to 12 foot intervals. Maintain a constant supply of fresh bait for 15 days or until signs of mouse activity cease.

Rats: Apply 6 to 23 blocks per placement. Space placements at 15 to 30 foot intervals. Maintain a constant supply of fresh bait for 10 days or until signs of rat activity cease.

Follow-up: Replace contaminated or spoiled bait immediately. Collect and dispose of all visible dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

(06004)